2018 OCO - GOSAT Technical Interchange Meeting at AGU

OCO-2 / OCO-3 Status

Minutes of the TIM

David Crisp and Annmarie Eldering

Jet Propulsion Laboratory, California Institute of Technology

for the OCO-2 Science Team

December 11, 2018




OCO-2 Status Summary

- Observatory Status: Nominal
 - Over 4.3 years and 23650 orbits completed
 - Last debris avoidance RMM (#7) on 11/05/2018
 - Next Drag Makeup Maneuver (DMM) tentatively scheduled for 10 March 2019 to coincide with an Inclination Angle Maneuver (IAM)
- Instrument Status: Nominal
 - Suspected Single Event Upset of SCO2 Data Processing Unit on 12/03/2018. DPU reset normally. 41 hours of data (orbits 23526-23550) lost
 - Error in slow degradation correction identified and will be corrected in B10
- Science Status: Nominal
 - 265 million soundings (through October 31, 2018) in the V8r collection publically available at GES DISC
 - V9r products through October 2018 publicly available at the GES DISC.
 - "Build 10" testing plan beginning to come together
 - ACOS/GOSAT version 9, slowly coming together
 - Production schedule is under review





Formation Flying with CALIPSO




OCO-2 maintained a tight formation with CALIPSO between May 2015 and September 2018, when CALIPSO left the A-Train.





Throughput Trending



An error in the slow degradation correction (Blue line in inset) introduced a small (~1%) radiometric error in the ABO2 throughput in the v8 product.






The OCO-2 V9 Product

- The OCO-2 Team released the Version 9 (V9) product on 10/15.
 - refined pointing
 - a correction to the prior meteorology
 - updated filtering and bias correction
- These updates
 - reduce bias in the presence of rough topography
 - Provide better sampling over topical and boreal forests with slightly more scatter
- This new dataset is available through the GES-DISC

https://disc.gsfc.nasa.gov/datasets? keywords=oco-2&page=1












Persistent X_{CO2} Anomalies Comparison of the V8 and V9 Products



 OCO-2 XCO₂ estimates are being used to quantify persistent anomalies associated with CO₂ emissions (sources) and uptake (sinks).

[Hakkarainen et al. 2018]










Persistent X_{CO2} Anomalies Comparison of the V8 and V9 Products



• While the V8 and V9 anomalies are similar, the V9 product has much less scatter in areas with rough topography (i.e. Himalayas, Canadian Rockies)


[Hakkarainen et al. 2018]







Improved Yields



The pointing correction, combined with re-tuned quality filters improved the yield, especially in regions with rough topography and over dark forested surfaces.







Subsetting Capability Added for OCO-2 "Lite" Files

- OCO-2 and ACOS Level 2 "Lite" files can now be subsetted spatially and by variable
 - Spatial subsets may be selected within a bounding box or within a user-defined radius around a user-specified location ("point+radius subsetting")
- This service is now operational https://disc.gsfc.nasa.gov/information/news?title=Subsetting%20capability%20added%20for%20OCO-2%20%22Lite%22%20files

Questions:

Dana Ostrenga, Thomas Hearty,
 Paul Huwe, Jennifer Adams,
 Andrey Savtchenko, Jerome
 Alfred, Lena Iredell





Left: XCO2 for two OCO-2 orbits from 13 (left) and 18 Aug (right) that passed within 100 km of the Carr Fire.







OCO-3 Summary

- OCO-3 shipped to Cape Canaveral on November 30
 - Almost finished with the check out and preparations for launch.
 - It will be stored until February, and will then go to Dragonland for integration in the Dragon trunk and then loading on the Falcon 9 launch vehicle.
 - Current launch date: March 16, 2019 from the Space-X launch pad





